arXiv:1703.02436v1 [math.GR] 7 Mar 2017

ADDENDUM TO SHARPLY 2-TRANSITIVE GROUPS OF CHARACTERISTIC 0

MALTE SCHERFF KATRIN TENT

Abstract. In this short note we show how to modify the construction of non-split sharply 2-transitive groups of characteristic 0 given in [RT] to allow for arbitrary fields of characteristic 0.

1. Introduction

The first sharply non-split sharply 2-transitive groups in characteristic 0 were constructed in [RT]. However, the construction given there only works when starting from the group $AGL(1,\mathbb{Q}) = \mathbb{Q}_+ \rtimes \mathbb{Q}^*$. We modify the construction given there in order to prove:

Theorem 1.1. *For any field $\mathbb{K}$ of characteristic 0 the group $AGL(1,\mathbb{K}) = \mathbb{K}_+ \rtimes \mathbb{K}^*$ can be embedded into a sharply 2-transitive group of characteristic 0 not containing any regular normal subgroup.*

To prove Theorem 1.1 we introduce the following equivalence relation (replacing the equivalence relation given in [RT]): for any group G and involution $j \in G$ we say that involutions $s, t \in G$ are *equivalent relative to j* (and write $s \approx_j t$) if Cen(js) = Cen(jt).

The following proposition replaces Proposition 1.3 of [RT] and provides the induction step for the proof of Theorem 1.1

Proposition 1.2. *Let G be a group containing involutions j, t and t' with $j^{t'} = t$ and $A = \mathrm{Cen}_G(j)$. Assume that G, j, t, t' and A satisfy assumptions (1) – (3) of Theorem 1.1. in* [RT] *and furthermore:*

- (4') *for any involution s with $s \approx_j t$ there is some $a \in A$ such that $s = t^a$.*
- (5') *for any involution $s \neq j$, we have $\mathrm{Cen}(js) = \{1\} \cup \{js' \colon s' \approx_j s\}$.*
- (6') *for any involution $s \notin AtA, s \neq j$, there is an involution $s' \in G$ with $s' \approx_j s$ such that $\mathrm{Cen}(js) = \langle js' \rangle$.*

Then for any involution $v \in G$ with $v \neq j$ there exists an extension G_1 of G such that for $A_1 = \mathrm{Cen}_{G_1}(j)$ there exists some $f \in A_1$ with $t^f = v$ and conditions $(1)-(3)$ and $(4')-(6')$ continue to hold with G_1 and A_1 in place of G, A.

Date: March 8, 2017.

2010 *Mathematics Subject Classification.* Primary: 20B22.

Key words and phrases. sharply 2-transitive, free product, HNN extension, malnormal.

Note that for the group $AGL(1,\mathbb{Q})$ the new equivalence relation agrees with the one given in [RT]. It is easy to see exactly as in [RT] that for any field $\mathbb{K}$ of characteristic 0, the group $AGL(1,\mathbb{K})$ satisfies properties (1) – (3) and (4') – (6').

Using the following lemma, the proof of Proposition 1.3 in [RT] carries over verbatim to this setting.

Lemma 1.3. *In the situation of Proposition 1.2 we have* $\mathrm{Cen}(js) = \mathrm{Cen}((js)^n)$ *for any involution* $s \in G$. *In particular, for involutions* $s, s' \in G$ *and* $n, m \in \mathbb{Z}$ *such that* $(js)^n = (js')^m$ *we have* $s \approx_j s'$.

Proof. Since js centralizes $(js)^n$, assumption (5') implies $s \approx_j j(js)^n$. The second part follows directly from this. □

As in [RT] we also see that for any field $\mathbb{K}$ of characteristic 0, the group $AGL(1,\mathbb{K}) * \mathbb{Z}$ satisfies properties (1) – (3) and (4') – (6'). Now the proof of Theorem 1.1 follows exactly as in [RT].

References

[Ne] B. H. Neumann, *On the commutativity of addition*, J. London Math Soc. 15 (1940), 203–208.

[RT] E. Rips, K. Tent, *Sharply* 2*-transitive groups of characteristic* 0, to appear in J. Reine Angew. Math.

Malte Scherff, Katrin Tent, Mathematisches Institut, Universität Münster, Einsteinstrasse 62, 48149 Münster, Germany

E-mail address: tent@wwu.de